Galore International Journal of Health Sciences and Research
[Indexed, Peer-Reviewed Journal]
Vol. 11; Issue: 4; Oct.-Dec. 2026
Website: www.gijhsr.com
Review Article *P-ISSN: 2456-9321*

Glycerine as an Alternative Extraction Vehicle in Homoeopathic Mother Tincture Preparation: A Review

Dr. G. Rathna Kumari[1], Dr. Rangala Sai Siva Harshini[2]

[1]Professor and HOD, Department of Homoeopathic Pharmacy, MNR Homoeopathic Medical College, Sangareddy, Telangana, India.
[2]Postgraduate Scholar, Department of Homoeopathic Pharmacy, MNR Homoeopathic Medical College, Sangareddy, Telangana, India.

Corresponding Author: Dr. Rangala Sai Siva Harshini

DOI: https://doi.org/10.52403/gijhsr.20260416

ABSTRACT

Alcohol is conventionally used as an extraction vehicle in homoeopathic mother tincture preparation because of its solvent and preservative properties. However, interest in alternative extraction vehicles has encouraged the exploration of glycerine owing to its distinctive physicochemical characteristics. This review examines the potential of glycerine as an alternative extraction vehicle in Homoeopathic Pharmacy, focusing on its extraction mechanisms, solubilization of phytoconstituents, advantages, and limitations. Glycerine's hydrogen-bonding capacity, hygroscopicity, and low volatility may support the extraction of selected plant constituents and the development of ethanol-free liquid preparations. Nevertheless, its high viscosity, selective extraction characteristics, analytical challenges, and uncertain microbial stability require careful consideration. The suitability of glycerine depends on the medicinal plant, solvent composition, and preparation conditions. Further plant-specific comparative studies evaluating phytochemical profiles, extraction yield, physicochemical properties, microbial quality, and stability are necessary to establish its pharmaceutical applicability. Glycerine should therefore be considered a potential alternative extraction vehicle requiring scientific validation rather than an established replacement for alcohol in homoeopathic mother tincture preparation.

Keywords: Glycerine, Homoeopathic mother tincture, Extraction vehicle, Herbal extraction, Pharmaceutical standardization, Homoeopathic pharmacy.

INTRODUCTION

Medicinal plants continue to play a pivotal role in global healthcare owing to their rich reservoir of bioactive compounds and diverse therapeutic applications. Despite remarkable advances in synthetic pharmaceuticals, plant-derived medicines remain integral to traditional and complementary healthcare systems and continue to contribute significantly to modern drug discovery. The increasing demand for safe, effective, and evidence-based herbal formulations has emphasized the importance of scientific validation, quality assurance, and standardization of medicinal plant preparations. [1]

Medicinal plants have served as an indispensable source of healthcare since ancient times and continue to play a vital role in the prevention and management of various diseases. Owing to their rich diversity of bioactive phytoconstituents, medicinal plants remain integral to traditional, complementary, and modern healthcare systems, contributing significantly to drug discovery and therapeutic innovation. Despite remarkable advances in synthetic pharmaceuticals, there is a growing global interest in plant-derived medicines because of their therapeutic efficacy, accessibility, and potential for reduced adverse effects. Consequently, increasing emphasis has been placed on the scientific validation, quality assurance, and standardization of herbal preparations to ensure their safety, efficacy, and consistency. [1,2]

The selection of an appropriate extraction vehicle is a fundamental consideration in the preparation of herbal pharmaceutical formulations, as it directly influences the extraction efficiency, chemical composition, stability, and overall quality of the final product. An ideal extraction vehicle should possess suitable solvent properties, chemical stability, safety, and compatibility with the phytoconstituents present in medicinal plants. Various solvents, including water, alcohol, hydroalcoholic mixtures, glycerine, and other pharmaceutical vehicles, are employed depending on the nature of the plant material and the desired therapeutic constituents. Among these, alcohol has traditionally remained the principal extraction vehicle in homoeopathic mother tincture preparation because of its excellent solvent capacity and preservative properties. However, the increasing demand for safer and alcohol-free pharmaceutical alternatives has encouraged scientific interest in exploring glycerine as a potential extraction vehicle.[3,4]

KEY ADVANTAGES OF USING GLYCERINE

1. **Enzyme Stabilization:** Glycerin is a well-established cryoprotectant that prevents the formation of damaging ice crystals when enzyme solutions are stored at low temperatures, such as -20°C. This significantly enhances the long-term stability and shelf-life of enzymes. Many commercially available enzymes, like restriction enzymes and polymerases, are supplied in buffers containing 50% glycerol.
2. **Enhanced Solubility:** In pharmaceutical formulations, glycerin can act as a co-solvent to increase the solubility of enzyme-based drugs, which is particularly useful for drug delivery systems.
3. **Modulation of Enzyme Kinetics:** The increased viscosity of glycerin-containing solutions can slow down enzymatic reaction rates. This property can be advantageous in certain experimental setups, such as stopped-flow kinetics, where precise control over reaction times is crucial.
4. **Compatibility with Analytical Techniques:** Glycerin generally does not interfere with common analytical methods like UV-Vis spectroscopy, circular dichroism, or fluorescence, making it a suitable component in various assay buffers.[5,6]

MATERIALS & METHODS

This article is based on information compiled from standard homoeopathic textbooks, and relevant articles and websites.

PHYTOCHEMICAL EXTRACTION PROFILE OF GLYCERINE

Glycerine is a highly polar extraction vehicle that exhibits excellent affinity for hydrophilic and moderately polar phytoconstituents, including phenolic compounds, flavonoids, tannins, glycosides, saponins, polysaccharides, and certain alkaloids, while showing relatively lower

efficiency for highly lipophilic constituents such as essential oils and resins[7]

TABLE 1: Phytochemical extraction profile of glycerine

Well extracted by glycerine	Poorly extracted by glycerine
Phenolic compounds	Essential oils
Flavonoids	Waxes
Tannins	Resins
Glycoside	Non-polar terpenes
Saponins	Lipids/fats
Polysaccharides	Highly lipophilic compounds
Some polar alkaloids	—

MECHANISM OF GLYCERINE-ASSISTED EXTRACTION

Glycerine functions as an effective extraction vehicle owing to its high polarity, strong hydrogen-bonding ability, and hygroscopic characteristics, all of which contribute to the efficient recovery of bioactive constituents from medicinal plants. During maceration, glycerine gradually permeates the plant tissues, facilitating the diffusion of intracellular phytochemicals into the extraction medium. The presence of three hydroxyl (–OH) groups enables glycerine to interact with hydrophilic and moderately polar phytoconstituents, including phenolic compounds, flavonoids, tannins, glycosides, polysaccharides, and certain alkaloids, thereby enhancing their dissolution and extraction. In addition, its hygroscopic nature helps maintain adequate moisture within the plant matrix, promoting continuous extraction while protecting sensitive phytochemicals from degradation. These combined properties make glycerine not only an efficient extraction vehicle but also a suitable medium for preserving the stability of therapeutically valuable plant constituents. [4,7]

INTERACTION OF GLYCERINE WITH PLANT MATRIX

The initial stage of extraction involves the contact of the extraction vehicle with the structural components of the medicinal plant. During maceration, glycerine comes into contact with the plant surface and wets the material, allowing the vehicle to gradually access the porous and intercellular regions of the plant matrix. This interaction is facilitated when the crude drug is appropriately reduced in size, as cutting, crushing or powdering increases the surface area available for contact with the extraction medium. The accessibility of glycerine to the plant matrix may also vary according to the anatomical structure of the drug, its moisture content and the composition of the extraction vehicle.[8,4]

Once adequate contact is established, hydration of the plant material can increase the accessibility of soluble intracellular constituents to the surrounding vehicle. The hydroxyl groups present in glycerine allow it to interact with water molecules and polar functional groups of plant constituents through hydrogen bonding. Such interactions can favour the dissolution of compatible phytochemicals after their release from the plant matrix. [3] As the concentration of extractable constituents becomes higher within the plant material than in the surrounding vehicle, a concentration gradient is established, facilitating their movement from the plant matrix into the extraction medium.[7,8]

Thus, the interaction between glycerine and medicinal plant material is a progressive process involving surface wetting, hydration, movement through the plant matrix, release of soluble constituents and their subsequent diffusion into the extraction vehicle. The overall efficiency of extraction is determined by the combined influence of glycerine's physicochemical characteristics and the properties of the plant material, together with the conditions maintained during maceration.

DIFFUSION AND MASS TRANSFER DURING GLYCERINE EXTRACTION

Following contact with the plant matrix, extraction involves the transfer of soluble constituents from the plant material into the surrounding extraction medium. This transfer is influenced by the concentration gradient between the plant matrix and the extraction solvent and is affected by factors such as solvent properties, particle size and contact time.[1,2] Glycerine can solubilize compatible constituents released from the plant matrix, allowing their gradual movement into the extraction medium.[3] Continued contact during maceration facilitates progressive extraction of soluble constituents, contributing to their accumulation in the resulting glycerine extract.[9]

SOLUBILIZATION OF PHYTO CONSTITUENTS BY GLYCERINE

Glycerine has a strong affinity for polar and water-soluble plant constituents, which contributes to its extraction potential. Its three hydroxyl groups promote hydrogen-bonding interactions with compatible phytochemicals and facilitate their dissolution in the extraction medium.[10,11] Phenolic compounds, flavonoids, tannins, glycosides and polysaccharides may therefore be extracted to varying degrees according to their chemical properties and the composition of the extraction system. The addition of water can alter the polarity of the glycerine-based medium and improve the solubilization of constituents with different polarities. Hence, the proportion of glycerine and water can influence both the nature and extent of phytoconstituents extracted during maceration.[7]

FACTORS INFLUENCING GLYCERINE-BASED EXTRACTION

The efficiency of glycerine-based extraction depends on several factors, including the nature of the plant material, particle size, glycerine-to-water ratio, extraction time, temperature, and the method of extraction. These variables influence solvent penetration, solubilization, and the transfer of phytoconstituents from the plant matrix into the extraction medium. [7,9]

Nature and particle size of the plant material: The chemical composition, anatomical structure, moisture content, and degree of comminution of the crude drug influence extraction efficiency. Reducing particle size increases the surface area available for contact with the extraction vehicle, potentially facilitating the release of soluble constituents. [4,8]

Glycerine-to-water ratio: The proportion of glycerine and water affects the polarity, viscosity, and solvent properties of the extraction medium. Water may improve the hydration of plant tissues and facilitate the extraction of certain water-soluble constituents, whereas glycerine contributes to the solubilization of compatible polar compounds. Therefore, the optimal ratio may vary according to the medicinal plant and its phytochemical composition. [7,10]

Extraction time and temperature: Prolonged contact between the plant material and the extraction medium may promote the transfer of soluble constituents until equilibrium or practical extraction limits are approached. Temperature can influence diffusion, viscosity, and solubility; however, excessive heating may adversely affect heat-sensitive constituents. Appropriate extraction conditions should therefore be selected according to the stability of the target phytochemicals. [4,9]

Method of extraction: Maceration, agitation, and other extraction techniques differ in their ability to facilitate contact between the plant material and the solvent. During maceration, adequate contact time and periodic mixing may improve the uniformity of solvent exposure. The selection of a suitable method should consider the physical characteristics of the crude drug and the properties of the glycerine-based medium. [4,8]

Overall, these factors should be optimized and controlled to obtain reproducible glycerine-based extracts. Comparative evaluation of extraction yield, physicochemical characteristics, phytochemical profiles, and stability is necessary to determine the suitability of glycerine for individual medicinal plants.

GLYCERINE IN HOMOEOPATHIC MOTHER TINCTURE PREPARATION

In Homoeopathic Pharmacy, mother tinctures are fundamental starting preparations for the preparation of subsequent potencies, and their quality depends on the crude drug, extraction vehicle, method of preparation, and compliance with pharmacopoeial standards. Alcohol is conventionally employed for its solvent and preservative properties. However, the exploration of alternative vehicles has created interest in glycerine because of its solvent capacity, low volatility, and humectant properties. Its potential application in mother tincture preparation depends on its ability to extract relevant phytoconstituents while maintaining acceptable physicochemical quality and stability. Therefore, systematic evaluation of extraction yield, physicochemical parameters, phytochemical composition, and microbial quality is essential to determine its suitability for homoeopathic pharmaceutical preparations. [4,7,10]

ADVANTAGES OF GLYCERINE-BASED MOTHER TINCTURES

1. Low volatility: Glycerine has low volatility compared with ethanol, which may reduce solvent loss during preparation and storage. [10]

2. Solvent properties: Its hydroxyl groups facilitate hydrogen bonding and the solubilisation of selected polar phytoconstituents. [10,11]

3. Reduced ethanol exposure: Glycerine provides a potential option for preparing liquid extracts without ethanol as the extraction vehicle. [10]

4. Hygroscopic nature: Glycerine absorbs moisture, which may influence the extraction characteristics of moist medicinal plant materials. [10]

5. Palatability: Its sweet taste may improve the acceptability of certain liquid preparations. [10]

6. Potential preservation benefits: Concentrated glycerine can reduce water availability, although the preservative effect depends on its concentration and the formulation. [10]

7. Alternative extraction potential: Glycerine may be investigated as an alternative extraction vehicle for selected plant constituents, depending on their chemical properties. [4,7,9]

DISADVANTAGES OF GLYCERINE-BASED MOTHER TINCTURES

1. High viscosity: Glycerine's viscosity may slow solvent penetration and mass transfer during extraction. [10]

Resolution: Optimize the glycerine-to-water ratio and consider controlled mixing to improve contact between the extraction vehicle and plant material. Any dilution should be evaluated for its effect on extraction efficiency and stability

2. Selective extraction: Glycerine may not extract all phytoconstituents as efficiently as appropriately selected hydroalcoholic solvents. [4,7,9]

Resolution: Glycerine is worth investigating because a different extraction vehicle may yield a different and potentially useful phytochemical profile.

3. Interference with total-solids estimation: Residual glycerine may contribute to the measured residue, requiring vehicle-blank correction when estimating extracted plant solids. [10]

Resolution: Analyze a corresponding vehicle blank under identical conditions and subtract the residual vehicle contribution to estimate the plant-derived solids more accurately.

4. Moisture-related variability: Its hygroscopic nature may affect the water

content and composition of the preparation during storage. [10]

Resolution: Standardize the initial moisture content of the crude drug, the glycerine-to-water ratio, and storage conditions. These measures can improve batch-to-batch consistency.

5. Handling difficulties: Its thick consistency may complicate filtration, transfer, dispensing and laboratory analysis. [10]

Resolution: These challenges may be minimized by using suitable filter media, vacuum-assisted filtration where appropriate, wide-bore or positive-displacement pipettes, and calibrated dispensers designed for viscous liquids. For analytical evaluation, validated sample dilution, vehicle-blank correction, and suitable analytical techniques, such as UV–Visible spectroscopy or high-performance liquid chromatography (HPLC), may improve measurement accuracy and reproducibility.

DISCUSSION

Glycerine offers potential as an alternative extraction vehicle in Homoeopathic Pharmacy owing to its distinctive solvent properties, hydrogen-bonding capacity, hygroscopicity, and low volatility. Its ability to extract selected phytoconstituents warrants investigation, as extraction efficiency depends on the chemical nature of the constituents and the medicinal plant. However, high viscosity, selective extraction, analytical challenges, and uncertainties regarding microbial stability require careful consideration. These limitations may be addressed through optimized extraction conditions, appropriate analytical methods, and systematic quality-control procedures. Comparative studies should evaluate phytochemical composition, extraction yield, physicochemical properties, and stability to determine the suitability of glycerine for individual medicinal plants. Therefore, glycerine should be investigated as a potential alternative rather than presumed to be equivalent or superior to alcohol in homoeopathic mother tincture preparation.

CONCLUSION

Glycerine offers a potential alternative extraction vehicle for homoeopathic mother tincture preparation, particularly for investigating the extraction of selected phytoconstituents from medicinal plants. This review highlights its pharmaceutical advantages, including low volatility, hygroscopicity, hydrogen-bonding capacity and suitability for developing ethanol-free liquid preparations, while recognising limitations associated with high viscosity, selective extraction and analytical standardisation. In the context of Homoeopathic Pharmacy, the use of glycerine warrants further investigation to determine its suitability for different medicinal plant materials and extraction procedures. Comparative experimental studies assessing physicochemical parameters, phytochemical composition, extraction yield, microbial stability and reproducibility are essential to establish its practical applicability. Therefore, glycerine should be considered a promising research avenue rather than an established replacement for alcohol. Evidence generated through systematic evaluation may contribute to the development of standardised alternative extraction methods for homoeopathic mother tincture preparation.

Declaration by Authors

Ethical Approval: Approved

Acknowledgement: None

Source of Funding: None

Conflict of Interest: The authors declare no conflict of interest.

Creative Commons Attribution License 4.0

This is an open-access article published under the terms of the Creative Commons Attribution 4.0 International License (CC BY 4.0).

https://creativecommons.org/licenses/by/4.0

REFERENCES

1. Latif R, Nawaz T. Medicinal plants and human health: a comprehensive review of bioactive compounds, therapeutic effects, and applications. Phytochem Rev. 2026; 25:2299-2342. doi:10.1007/s11101-025-10194-7.
2. World Health Organization. Monitoring health for the SDGs: the Global Health Observatory [Internet]. Geneva: World Health Organization; [cited 2026 May 13]. Available from: https://www.who.int/data/gho/data/themes/world-health-statistics
3. Homoeopathic Pharmacopoeia of India. Vol. I. New Delhi: Ministry of AYUSH, Government of India;
4. Azmir J, Zaidul ISM, Rahman MM, Sharif KM, Mohamed A, Sahena F, et al. Techniques for extraction of bioactive compounds from plant materials: a review. J Food Eng. 2013;117(4):426-436. doi: 10.1016/j.jfoodeng.2013.01.014.
5. Gekko K, Timasheff SN. Mechanism of protein stabilization by glycerol: preferential hydration in glycerol-water mixtures. Biochemistry. 1981;20(16):4667-4676. doi:10.1021/bi00519a023.
6. Vagenende V, Yap MG, Trout BL. Mechanisms of protein stabilization and prevention of protein aggregation by glycerol. Biochemistry. 2009;48(46):11084-11096. doi:10.1021/bi900649t.
7. Zhang QW, Lin LG, Ye WC. Techniques for extraction and isolation of natural products: a comprehensive review. Chin Med. 2018; 13:20. doi:10.1186/s13020-018-0177-x.
8. Chemat F, Vian MA, Cravotto G. Green extraction of natural products: concept and principles. Int J Mol Sci. 2012;13(7):8615-8627. doi:10.3390/ijms13078615.
9. Sarker SD, Nahar L. An introduction to natural products isolation. In: Sarker SD, Nahar L, editors. Natural Products Isolation. 3rd ed. Totowa (NJ): Humana Press; 2012. p. 1-25. doi:10.1007/978-1-61779-624-1_1.
10. Sheskey PJ, Cook WG, Cable CG, editors. Handbook of Pharmaceutical Excipients. 9th ed. London: Pharmaceutical Press; 2020.
11. Dai J, Mumper RJ. Plant phenolics: extraction, analysis and their antioxidant and anticancer properties. Molecules. 2010;15(10):7313-7352. doi:10.3390/molecules15107313.

How to cite this article: Dr. G. Rathna Kumari, Dr. Rangala Sai Siva Harshini. Glycerine as an alternative extraction vehicle in homoeopathic mother tincture preparation: a review. *Gal Int J Health Sci Res*. 2026; 11(4): 138-144.
DOI: https://doi.org/10.52403/gijhsr.20260416

******